

Your guide to your medication:

Medications Used in the Treatment of Dementia

The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions.

National University Hospital
5 Lower Kent Ridge Road
Singapore 119074
Tel: (65) 6779 5555 Fax: (65) 6779 5678
Website: www.nuh.com.sg
Company Registration No. 198500843R

Information is correct at time of printing (December 2016) and subject to revision without prior notice.

- Donepezil
- Rivastigmine
- Galantamine

Medicines used for the treatment of dementia

- The class of medicines commonly used to treat dementia is known as acetylcholinesterase inhibitors.
- These include:
 - **Donepezil** tablet
 - **Rivastigmine** capsule or transdermal patch
 - **Galantamine** capsule
- These medicines do not cure the disease but may delay the worsening of some symptoms of dementia for a period of time.

How these medications are administered

- Follow the directions on your medicine label or take as directed by your doctor.
- Oral medications are to be taken after meals.
- The Rivastigmine patch is meant to be applied on the skin. Please refer to the detailed product information available.

Possible common side effects (may affect up to 10% of patients)

Similar to all medications, the medications used for the treatment of dementia may cause some possible side effects in some people. Not everyone will experience the side effects as every individual responds differently to the medications. Inform your doctor or pharmacist should these side effects become too bothersome.

- Abdominal pain
- Diarrhoea
- Dizziness
- Headache
- Nausea, vomiting or loss of appetite
- Sleep disturbances
- Tiredness

For urgent medical attention

Consult your doctor as soon as possible if you have any of the following symptoms:

- Allergic reaction: skin rash or itch, difficulty in breathing, chest tightness, or swelling of the face, lips or other parts of the body
- Severe pain in the stomach
- Feeling faint
- Slow or irregular heartbeat

If you have any concerns about your medication, please consult your doctor or pharmacist.