

PATIENT INFORMATION LEAFLET

NON-STEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDs)

Diclofenac
Ibuprofen
Indomethacin
Ketoprofen
Mefenamic Acid
Naproxen
Piroxicam
Meloxicam
Celecoxib
Etoricoxib

ABOUT your medicine

This medicine can be used to treat pain and inflammation (swelling). It can also be used in rheumatoid arthritis, osteoarthritis and other inflammatory disorders.

The medicine works by blocking the production of prostaglandins. Prostaglandins are chemicals produced by the body after an injury, and they may cause pain and swelling. Reducing prostaglandin production will reduce symptoms after an injury. However, these prostaglandins also provide protection against gastric problems. Thus, NSAIDs may cause side-effects such as indigestion, ulcers, and stomach bleeding.

The newer types of NSAIDs, called the COX-2 selective NSAIDs (e.g. celecoxib, etoricoxib), selectively block the “inflammatory” prostaglandins but allow the stomach-protecting prostaglandins to remain in action. Gastric side-effects are therefore reduced.

HOW should I take the medicine?

Your doctor will determine the amount of medicine you should take based on the severity of your condition. Take this medicine in the dose prescribed by your doctor.

This medicine may be taken with food or directly after a meal to reduce gastric side-effects. COX-2 selective NSAIDs may be taken with or without food.

What should I do if **I FORGET** a dose?

If you miss a dose or forget to take your medicine, take it as soon as you remember. However, if it is almost time for your next dose, wait until

then to take the medicine and skip the missed dose. Do not double your dose or take extra medicine to make up for a missed dose. If you have missed several doses, please call your physician to check what you should do.

What are the possible **SIDE-EFFECTS** of the medicine?

The following are some of the more common side-effects:

- Gastro-intestinal discomfort
- Allergic reactions
- Nausea
- Diarrhoea
- Stomach ulcers
- Headache
- Dizziness
- Hearing disturbances.

Some people may experience fluid retention and raised blood pressure. NSAIDs may also affect blood clotting. Use of NSAIDs may also worsen asthma. Inform your doctor if you experience any side-effects.

What **PRECAUTIONS** should I take?

Before taking your medicine, inform your doctor if :

- you are allergic to any medicines.
- you are pregnant or planning to become pregnant.
- you are breast-feeding.

Keep all medication out of reach of children

- you have a history of gastric (stomach) or duodenal (intestinal) ulcers.
- you are on blood thinning agents (e.g. aspirin, warfarin).
- you are suffering from a defect of the blood clotting system.
- you have kidney, heart or liver problems.

How should I **STORE** the medicine?

Store the medicine in a dry place at room temperature, away from heat and direct light. Do not refrigerate your medicines unless instructed to do so.

National University Hospital

5 Lower Kent Ridge Road, Singapore 119074
Tel: (65) 6779 5555 Fax: (65) 6779 5678
Website: www.nuh.com.sg

Company Registration Number. 198500843R

The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions.

Information is correct at time of printing and subject to revision without prior notice.

Discard all medication that has expired or is no longer required