


Posterior Tibial Tendon Dysfunction

Posterior Tibial Tendon Dysfunction (PTTD), also known as Adult Acquired Flatfoot Deformity, is a condition caused by overuse of the posterior tibial tendon. Inflammation and tears in the tendon make it weaker and less able to support the foot, resulting in pain and swelling.

If left untreated, it may also result in a permanent flattening of the arch.



Common risk factors of PTTD:



Common treatment options:



Activity modification – Reduce high impact activities such as running and jumping



Cold compress – To reduce swelling and provide relief. Put on a sock and place an ice pack over the affected area for 10 to 15 minutes



Offloading devices – An ankle brace and/or a removable cast walker can offload pressure from the injured tendon



Orthoses – For better control of foot function and to reduce strain on your tendon



Footwear – Choose shoes with appropriate support or cushioning to reduce strain on the tendon



Stretches – Calf stretches help restore necessary movement in the lower limb

Other treatment options:



Strengthening exercises for the lower limb



Surgery

For more information, kindly consult your podiatrist.

Scan the QR codes for more information:



**Stretches and Exercises
for PTTD**



**How to Choose
Appropriate Footwear**



FAQs on PTTD

National University Hospital
5 Lower Kent Ridge Road Singapore 119074
Tel: (65) 6908 2222
Website: www.nuh.com.sg

The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions.

Information is correct at time of printing (October 2021) and subject to revision without prior notice.

Copyright (2021). National University Hospital
All rights reserved. No part of this publication may be reproduced without permission in writing from the National University Hospital