

PATIENT INFORMATION LEAFLET

ANTI-FUNGAL EYE PREPARATIONS

Natamycin Eyedrops
Amphotericin Eyedrops

ABOUT your medicine

Anti-fungal eye preparations are used to treat some types of fungus infections of the eye. This medicine may be given alone or with other medicines that are taken by mouth for eye infections.

HOW should I use the medicine?

Your doctor will determine the amount of medicine you should use based on the severity of your condition. Use this medicine in the dose prescribed by your doctor.

Proper administration of eye preparations:

- Wash your hands with soap and water and read the instructions on the label.
- Shake the bottle before use.
- Tilt your head backwards and look up.
- Gently pull the lower eyelid down to form a pocket and squeeze the bottle to apply a single drop into it. To prevent contamination, please ensure that the tip of the bottle does not touch the eye or any other surface.
- For ointments, apply a 0.5-1cm ribbon to the inner side of the lower lid.
- Keep your eyelid closed for at least 30 seconds (do not blink, rub, wipe or squeeze the eye).
- Gently apply pressure to the inner corner of the lower eyelid with your finger for 1 or 2 minutes to allow the medicine to be absorbed by the eye.
- If you need to put in more than one drop, or if you are using another type of eyedrops, you should wait for 5 to 10 minutes before putting the next drop in. (This is to prevent the first drop from being washed out by the second before it has time to work, and to prevent interaction between 2 different types of eyedrops.)

Keep all medication out of reach of children

- If you are using eyedrops as well as eye ointments, you should use the eyedrops first.

If you are wearing contact lenses, do not wear them until your entire course of medicine is completed.

Eyedrops should not be used for longer than 4 weeks after opening. Write down the date that you opened the bottle on the label so that you will know when to throw it away.

What should I do if **I FORGET** a dose?

If you miss a dose or forget to use your medicine, use it as soon as you can. However, if it is almost time for your next dose, wait until then to use the medicine and skip the missed dose. Do not double your dose or use extra medicine to make up for a missed dose.

What are the possible **SIDE-EFFECTS** of the medicine?

Side-effects that may occur include eye irritation, redness or swelling, blurred vision, eye pain, and sensitivity to light. You may or may not experience any of these side-effects. These side-effects may go away during treatment as your body adjusts to the medicine. Check with your doctor if the effects persist.

Seek medical attention if any of the following side-effects occur:

- Rash or redness around the eyes
- Redness or swelling of the membrane covering the white part of the eye

Keep all medication out of reach of children

Discard all medication that has expired or is no longer required

What **PRECAUTIONS** should I take?

Inform your doctor if:

- you are allergic to any other medicines, or if you develop an allergy while taking this medicine. The symptoms of a drug allergy include one or more of the following: swollen face/eyes/lips, difficulty in breathing or widespread itchy skin rashes.
- your eye condition does not improve after 5 to 7 days or if it worsens.

How should **I STORE** the medicine?

Store the medicine in a dry place at room temperature, away from heat and direct light. Do not refrigerate your medicines unless instructed to do so.

National University Hospital

5 Lower Kent Ridge Road, Singapore 119074
Tel: (65) 6779 5555 Fax: (65) 6779 5678
Website: www.nuh.com.sg

Company Registration Number. 198500843R

The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions.

Information is correct at time of printing and subject to revision without prior notice.

Discard all medication that has expired or is no longer required