

ABOUT your medicine

The class of medicines commonly used to treat Alzheimer's disease is known as cholinesterase inhibitors. They are designed to prevent the breakdown of acetylcholine, a chemical messenger in the brain that is important for memory and other thinking skills. The medicine works to keep levels of this chemical messenger high, even while the cells that produce the messenger continue to become damaged or die. They work most effectively for people who are mildly to moderately affected by the disease.

PATIENT INFORMATION LEAFLET**MEDICATIONS FOR ALZHEIMER'S DISEASE**

Medicines in this class include:

- Donepezil
- Rivastigmine
- Galantamine

HOW should I take the medicine?

Donepezil

Donepezil is usually taken once a day in the evening just before bedtime.

Rivastigmine & Galatamine

Take these medicines exactly as your doctor prescribes. The dose may differ for different patients. It is usually taken with meals.

These medicines must be taken regularly every day over a period of time to be effective. It is common for your doctor to start you on a low dose and gradually increase as required.

E110R0

Keep all medication out of reach of children

Discard all medication that has expired or is no longer required

What should I do if **I FORGET** a dose?

Take this medicine exactly as directed by your doctor. If you miss a dose, take it as soon as you remember unless it is almost time for the next scheduled dose. In that case, skip the missed dose and take the next one as directed. Do not double the dose to make up for the missed dose.

What are the possible **SIDE-EFFECTS** of the medicine?

Nausea, vomiting, abdominal pain, loss of appetite and increased frequency of bowel movements might be expected with any of these medicines. Other side-effects include headache, dizziness, fatigue, malaise and anxiety. Some patients may exhibit an initial increase in agitation but this should subside after the first few weeks of therapy.

What **PRECAUTIONS** should I take?

These medications may cause drowsiness. Do not drive or operate machinery.

Do not take other medicines unless they have been discussed with your doctor or pharmacist especially non-prescription medicines such as painkillers, e.g. aspirin or ibuprofen and medications for cough and cold.

Tell your doctor about any nausea, vomiting, diarrhoea, stomach pain or loss of appetite.

It is very important that you see your doctor for regular visits and report worsening of your condition or if you notice any new symptoms.

How should **I STORE** the medicine?

Store the medicine in a dry place at room temperature, away from heat and direct light. Do not refrigerate your medicines unless instructed to do so.

National University Hospital

5 Lower Kent Ridge Road, Singapore 119074
Tel: (65) 6779 5555 Fax: (65) 6779 5678
Website: www.nuh.com.sg

Company Registration Number. 198500843R

The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions.

Information is correct at time of printing and subject to revision without prior notice.

Keep all medication out of reach of children

Discard all medication that has expired or is no longer required